

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

CR 167804

E83-10190

AgRISTARS

SR-B2-04375

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

A Joint Program for
Agriculture and
Resources Inventory
Surveys Through
Aerospace
Remote Sensing

Supporting Research

November 1982

FINAL REPORT

LABELING RESEARCH IN SUPPORT OF THROUGH-THE-SEASON AREA ESTIMATION

R. N. Colwell, Principal Investigator
C. M. Hay, E. J. Sheffner
University of California, Berkeley
Space Sciences Laboratory
Berkeley, California 94720



(E83-10190) LABELING RESEARCH IN SUPPORT OF
THROUGH-THE-SEASON AREA ESTIMATION Final
Report, 15 Nov. 1980 - 30 Jun. 1982
(California Univ.) 103 p HC AC6/MF A01

N83-20312

Unclass
00190

CSCI 02C G3/43



Earth Resources Research Division
Lyndon B. Johnson Space Center
Houston, Texas 77058

TECHNICAL REPORT STANDARD TITLE PAGE

1. Report No. SR-B2-04375		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle Labeling Research in Support of Through-the-Season Area Estimation				5. Report Date November 1982	
				6. Performing Organization Code	
7. Author(s) C.M. Hay and E.F. Sheffner				8. Performing Organization Report No. Series 23, Issue 10	
9. Performing Organization Name and Address Remote Sensing Research Program 260 Space Sciences Laboratory University of California Berkeley, California 94720				10. Work Unit No.	
				11. Contract or Grant No. NAS9-14565	
12. Sponsoring Agency Name and Address NASA - Johnson Space Center Houston, Texas 77058				13. Type of Report and Period Covered Final Report - 15 May 75 to 30 June 82	
				14. Sponsoring Agency Code	
15. Supplementary Notes					
16. Abstract This report documents work done to support the development of Landsat-based through-the-season labeling procedures for corn and soybeans. A model for predicting labeling accuracy within key time periods throughout the growing season is outlined. Two methods for establishing the starting point of one key time period, viz., early season, are described. In addition, spectral-temporal characteristics for separating crops in the early season time period are discussed.					
17. Key Words (Selected by Author(s)) Corn and soybeans, early season, Landsat, labeling accuracy				18. Distribution Statement	
19. Security Classif. (of this report) Unclassified		20. Security Classif. (of this page) Unclassified		21. No. of Pages 102	
22. Price*					

Space Sciences Laboratory
University of California
Berkeley, California 94720

Final Report

for

NASA Contract NAS9-14565

LABELING RESEARCH IN SUPPORT OF
THROUGH-THE-SEASON AREA ESTIMATION

Principal Investigator

Professor R.N. Colwell

Project Scientists

C.M. Hay (Project Manager)

E.J. Sheffner

Period Covered

15 November 1980 - 30 June 1982

Remote Sensing Research Program
Space Sciences Laboratory Series 23, Issue 10

LABELING RESEARCH IN SUPPORT OF THROUGH-THE-SEASON AREA ESTIMATION

**Claire M. Hay
Edwin J. Sheffner**

1.0 INTRODUCTION AND RESEARCH GOALS

The Large Area Crop Inventory Experiment (LACIE) demonstrated the usefulness of Landsat data for crop identification measurements in support of area estimation without a ground sample stage. In particular, LACIE area estimation procedures were for estimation of small grain acreages in areas where ground data would not be available. The LACIE small grains procedures were end-of-season procedures. That is, the LACIE procedures depended upon the full set of crop season acquisitions in order to perform the crop type labeling operation with sufficient accuracy. During the LACIE Transition Program, and in the subsequent AgRISTARS Project, there was a desire to not only extend the LACIE-like technology to crop types other than small grains, but also to more directly address the need for a continual through-the-season estimation capability. Through-the-season capability was most desirable to all potential users of the technology, since crop production indications prior to the end-of-season would allow for more effective management and planning within the agrobusiness community and commodities market.

The crop types selected for the extension of the LACIE-like technology were corn and soybeans. The particular foreign areas of interest for application of the technology were Argentina and Brazil. Therefore the work described in this report was directed toward corn and soybean crop types. The ultimate goal of the work was to develop corn and soybean crop type labeling procedures to be used in support of a through-the-season area estimation technology. It was envisioned that a through-the-season capability would be developed that would allow for the estimation of crop acreages at any time in the growing season. This may mean that more than one pre-end-of-season labeling procedure would need to be developed for use in different periods within the growing season. At the least, it was felt that the particular point in the season at which an estimate was desired would affect the operating parameters of any labeling procedure. Therefore, a further goal of this research was to come to understand the dynamics of crop separability as the growing season progressed.

2.0 GENERAL APPROACH

There were three phases planned in the overall approach to developing through-the-season labeling procedures. These three phases were:

First Phase: Determination and Description of Summer Crop Type Separation Characteristics Throughout the Growing Season

Second Phase: Development of Labeling Guidelines for Pre-End-of-Season Labeling of Corn and Soybeans

Third Phase: Development of Labeling Procedures to Support a Through-the-Season Area Estimation Procedure

Briefly in the first phase--Determination and Description of Summer Crop Type Separation Characteristics Throughout the Growing Season--the research would focus on determining what, if any, crop type or crop group characteristics could be observed within Landsat data that would be useful in uniquely identifying the crop types or at the very least the crop groups of interest. In addition, the expected improvement in labeling accuracy as the growing season progressed was to be evaluated and some mechanism for determining or for controlling the setting of the operating parameters at various periods in the growing season was to be investigated.

The second phase of research--Development of Labeling Guidelines--would formulate consistent labeling guidelines for the separation of corn and soybeans throughout the growing season. Clearly formulated guidelines would then allow labeling procedures to be designed in support of specific through-the-season area estimation procedures being developed.

The third phase--Development of Labeling Procedures--was to proceduralize the labeling guidelines into a consistent logical processing of the data either by an analyst, machine, or both interactively, similar to the proceduralization effort undertaken in the design of the Corn/Soybean Baseline Procedure C/S-1.

This document will report on the progress of the first phase of the research. At present no plans exist to continue with the other two phases due to termination of support for this effort.

3.0 THROUGH-THE-SEASON MODEL OF LABELING ACCURACY

3.1 GENERAL MODEL CONCEPT

There are four characteristics observable in multitemporal Landsat data that are most useful in determining the crop type or crop group identification of an agricultural feature. These characteristics are: (1) a temporal characteristic which pertains to the development of a feature's vegetation canopy through time, i.e., what time of year is a green vegetation canopy observable for the feature as opposed to bare ground; (2) a spectral characteristic which refers to the value of a given spectral variable, or to the relative relationship between two or more spectral variables for a given feature; and (3) two spatial characteristics, one which refers to the geometric attributes of a given feature such as the nature of a feature's boundary, and the other which refers to the areal distribution of similar and dissimilar features within the area of analysis. The most important characteristic for crop identification is the temporal characteristic. The temporal characteristic allows the separation of crop groups within a region, or more precisely, is used to define the various crop groups within the region. In order to support evaluation of the temporal characteristic, collateral data in the form of crop calendars are needed. Indeed, the most important piece of collateral data for crop type analysis is crop calendar data. The crop calendar data provides information about when crops within a given region are likely to be or are (if a real time crop calendar is used) in any given biostage.

From past research and experience, the spectral response of many vegetation biostages are familiar and generally known. Therefore a spectral crop calendar can be generated from phenological crop calendar data. This spectral crop calendar model allows an analyst or programmed decision algorithm to judge the most likely crop type or cover class label to assign to a given feature in a given set of multitemporal, spectral data such as Landsat data.

The logic process by which a labeling target's observed characteristics are evaluated can be illustrated by the list of "important questions" in Table 1. The ability to answer the important questions in a deterministic manner is dependent upon the temporal set of acquisitions available for analysis. It can be appreciated that as the crop growing season starts, there is little or no spectral data available to answer any of the questions. As the season progresses, however, more data become available and more of the "important questions" can be answered. The accuracy of any labeling analysis task is a function of the temporal sequence of acquisitions available. Thus, it can be understood that the labeling accuracy increases as a function of time. Note, however, that some questions can only be answered after a discrete event has occurred. For example, question 2a ("When does the maximum value for the green vegetation indicator occur?"): This question cannot be answered until some time AFTER the occurrence of the maximum value because a relative maximum can only be identified after the value of the green vegetation indicator has started to decrease again. This time delay in ability to answer a specific question relative to the occurrence of some events says that the labeling accuracy function relative to time is not a smoothly increasing function, but is a function which increases in steps. The plot of this function through the crop growing season would have several plateaus where the labeling accuracy would remain fairly constant until sufficient data had been received that would allow the answering of additional "important questions." (A graphical representation of this model is shown in Figure 1.) The above description of the model of labeling accuracy has ignored the question of data dropout and the use of multiyear labeling procedures or estimators. If multiyear labeling procedures or estimators were employed, the value for A_0 (initial value for labeling accuracy at the start (t_0) of a given crop year) would probably be higher (indicated by A_m). The labeling accuracy function would then proceed to perform generally as described above except for the increased accuracy levels expected from the use of multiyear data. Data dropout would have the effect of prolonging the length of various labeling accuracy plateaus until such time as sufficient data became available to answer additional "important questions."

With the above-described Labeling Accuracy Model as a basis, the Through-the-Season labeling research effort sought to accomplish the following objectives in Phase One:

- (1) Determine the specific identification characteristics that could be used to uniquely identify summer crop types (corn/soybean) or crop group to the highest accuracy level possible within any given accuracy plateau;
- (2) Determine a means of specifically identifying when each of the accuracy plateaus started and ended for the corn/soybean situation.

3.2 SPECIFIC THROUGH-THE-SEASON MODEL OF LABELING ACCURACY FOR CORN/SOYBEANS IN U.S. CORN BELT AND PERIPHERY

The accuracy plateaus described in the general Through-the-Season labeling model need to be more definitively described for the specific corn/soybean situation. This was done based on when each of the important questions from Table 1 could be answered during the Corn/Soybean Summer Crop growing season in the U.S. Corn Belt and Corn Belt Periphery. Figure 2 shows the specific characterization of the Labeling Accuracy Model for the Corn/Soybean Summer Crops in the Corn Belt Area. Use of multiyear data or data dropout were ignored for the present. The C/S-1 (Corn/Soybean Baseline Procedure) Summer Crop biowindows and the potential and actual separation windows have also been delineated on the specific model representation. As can be seen, C/S-1 is actually a mid-season procedure based on the occurrence of the actual separation window in the mid-season of the model. Since the DFS stratification in C/S-1 can be done with a minimum set of two acquisitions--one acquisition from the very early or pre-season time period and one acquisition from the very early to mid-season time period--C/S-1 can operate as a mid-season procedure. Partly because of the above reason and partly due to efficiency of approach, the decision was made to concentrate on the early season plateau in the initial efforts toward developing a Through-the-Season labeling logic.

Therefore the specific objectives of this initial Through-the-Season effort were:

- (1) to determine when t_2 occurred - for the Corn/Soybean situation,
 - (i) investigate how the occurrence of t_2 could be predicted operationally (i.e. real time during an actual Through-the-Season Estimation Effort)
- (2) to determine what characteristics or attributes of corn/soybeans, summer crops, and others were useful in separating
 - (i) summer crops from other (at the least)
 - (ii) corn and soybeans from other summer crops and from each other if possible.

4.0 DATA ANALYSIS METHODS

4.1 DATA SET

4.1.1 Segment Selection

The Through-the-Season labeling guidelines research was intended to support Through-the-Season area estimation procedures for use in foreign areas, particularly Argentina. However at the time of this study, no Argentina spectral data were available. For this reason the labeling guidelines research was conducted using the spectral data from segments located within the U.S. Corn Belt and Corn Belt Periphery. Segments selected for inclusion in the study were chosen

based on the following: (1) how well they represented the geographical diversity within the Corn Belt and Corn Belt Periphery, and (2) the adequacy of their Landsat acquisition histories. A good acquisition history was one that included at least one acquisition prior to the earliest planting date for corn, at least one acquisition during or just past corn tasseling or soybean pod setting, and a sufficient number of acquisitions in between these two bounding crop biostages. The dates of corn planting, corn tasseling, and soybean pod setting, as well as other crop calendar information was determined from the statewide year-specific crop calendars published by the U.S.D.A. - Statistical Reporting Service within their Weekly Crop and Weather Bulletins. Before final inclusion of a segment within the study data base, all segment acquisitions were visually screened for cloud cover, haze, and bad data lines. Eighteen segments were selected to comprise the data set. The list of selected segments and acquisitions is given in Table 2, and the segment locations are shown in Figure 3.

4.1.2 Segment Preprocessing

Sun angle and haze correction was performed on all acquisitions selected for inclusion in the data set. The sun angle and haze correction algorithm used to normalize the spectral data was the global XSTAR algorithm² developed by the Environmental Research Institute of Michigan (ERIM). This algorithm had been used quite successfully on numerous previously analyzed data sets of Landsat spectral data. Following the application of XSTAR, the spectral features of interest for the analysis were extracted. The spectral features extracted for analysis were as follows:

- (a) Tassel-cap component 1 (TC-1, Brightness)
- (b) Tassel-cap component 2 (TC-2, Greenness)--Green vegetation indicator
- (c) GRABS³ (Greenness Above Bare Soil)--Green vegetation indicator

Tassel-cap Brightness (TC-1) and GRABS were given the most emphasis in the analysis due to the successful results achieved in labeling crop type from prior studies.

Tassel-cap transformation is an affine transformation of the Landsat four-channel MSS data into four new channels or components of data which correspond more directly with physical processes and relationships, (particularly for vegetational ground features) than the landsat MSS bands do directly. GRABS is a green vegetation indicator (measure of infraredness) derived from tassel-cap greenness and the threshold of detection for green vegetation established for the 7/5VI:³

$$\text{GRABS} = (\text{TC-2}) - 0.09178 (\text{TC-1}) + 5.58959.$$

4.1.3 Sampling of Fields by Crop Type

Digitized ground data supplied by Johnson Space Center (JSC) was used to identify specific fields within a given crop type or vegetation cover class. A maximum of 10 fields for each summer crop type present within a segment was sampled. Sufficiently large non-summer crop fields were also sampled if they were present within a given segment. To minimize the effects of mis-registration between acquisitions, and to insure some level of spectral purity within the crop type samples, only field center pixels were included within the sample. A pixel was included in the sample of a field if it had a margin around it that was at least two pixels wide, or if the surrounding pixels were of the same cover class. This sampling strategy had the effect of excluding small fields from the study. In total 574 fields were sampled from the 18 segments in the data set. The number of fields sampled by segments and cover class is shown in Table 3.

4.1.4 Analysis Approach

For each of the sampled fields, a field mean was calculated in each of the spectral features to be analyzed for each acquisition. All of the analyses were then carried out on the field means data. Temporal plots of the field means were generated for Brightness (TC-1) and GRABS spectral features for each sampled field. In addition, segment level (segment average) temporal plots were generated for each segment by using a weighted (by pixel number) average of the field means within each crop type within each segment. Appendix A contains the segment mean temporal plots for Brightness and GRABS for each cover type sampled.

In order to determine whether there were any detectable differences between the crop groups, and the summer crop types at various points earlier in the growing season prior to end-of-season, the above described temporal plots were examined and evaluated with particular emphasis on observing differences as early in the growing season as possible. In addition the data were evaluated to try to determine at what points in the growing season significant increases in identification accuracies would likely occur as a function of the additional spectral and crop calendar data which become available as the season progressed.

5.0 DETERMINATION OF t_2 : SUMMER CROP SPECTRAL EMERGENCE DATE

5.1 DEFINITION, RATIONALE AND ASSUMPTIONS

In the Corn Belt, soybean is frequently the last summer crop planted. This is so partly because corn yield is more favorably affected by early planting than is soybeans. Thus corn planting has priority over soybeans; also soybeans can be double cropped after small grains so that double cropped soybean planting is delayed until after small grain harvest. Observations from past research of the temporal patterns of corn, soybean,

and other summer crops in the Corn Belt have shown this later planting pattern for soybeans to be reasonably consistent within the region, though local variations do occur. The date of t_2 is defined as the date when all summer crops are detectable. Within the Corn Belt t_2 can be approximately defined as the date when all soybeans are detectable. The word "all" in these definitions should be read $\geq 95\%$. This is similar to the manner in which the spectral biowindows were defined for C/S-1.

Since by definition from the model, the "early season" labeling guidelines would be based on the spectral-temporal characteristics of detectable vegetation, it was necessary to know or to be able to estimate when "all" summer crops were detectable in order to correctly operate the appropriate labeling procedures.* That is, the actual date of occurrence of t_2 needed to be determined. A method for estimating the occurrence of t_2 prior to its occurrence was also expected to be needed in the eventual Through-the-Season Estimation Procedure to be developed.

Since t_2 was expected to vary between years, a long-term average crop calendar would not be adequate for estimating t_2 as closely as was felt necessary. Therefore, a real-time method for estimating the occurrence of t_2 in any given inventory year was needed. The assumption was made that, at the least, some real-time crop calendar model data would be available. However, since no crop calendar model data was available for evaluation, and only limited progress on a Corn Crop Calendar Model was perceived at the time by these investigators, reliance upon such crop calendar model data was minimized as much as possible. The only year-specific crop calendar data available for this study, however, was the U.S.D.A.-S.R.S.'s Statewide Crop Calendars. Therefore the statewide year-specific crop calendar information was used to develop the method for estimating the occurrence of t_2 . It was hoped that crop calendar models under development could supply the needed information in foreign areas at some time in the not-too-distant future.

Since t_2 was expected to vary each year within each region, a reference point or start date t_0 was required that could be determined for each region each year. From the statewide year-specific U.S.D.A.-S.R.S. Crop Calendars t_0 was defined as the day on which 50% of the first planted summer crop was reported to have been planted. Thus in the Corn Belt, t_0 was defined as the date on which 50% of corn was reported to have been planted. From past research in crop calendar analysis for summer crops and small grains, it was determined that the 50% planted date for any crop tended to be less variable year-to-year than other percent planted points e.g. 5%, 75%, 95%, etc. Based on this perceived tendency toward higher stability, the 50% planted date was chosen as t_0 . In addition, this definition of t_0 allowed t_0 to be determined from data that would be available reasonably early in any growing season so that true Through-the-Season procedures could be developed that would start with the season and not at some later point in the season. (Past definitions of early and mid-season

* Labeling guidelines for a period $\leq t_2$ are called Very-Early Season guidelines by definition and may not rely on detectable vegetation for all potential summer crop fields.

have been stated in terms of days preceding harvest. One, therefore, needed an estimate of when harvest was to occur before one could classify acquisitions as to early or mid-season.)

The time between t_0 and t_2 (time interval $t_2 - t_0$) needed to be determined and its year-to-year variation evaluated.

5.2 ESTIMATION OF $(t_2 - t_0)$

Figure 4 graphically represents one preliminary method used to estimate $(t_2 - t_0)$. 1978 data from segment 127, Montgomery County, Indiana, was used to make a "first-look" sample calculation of the interval from 50% corn planted (t_0) to 95% soybean detectability (t_2). Segment 127 was selected from the 18 sampled segments because it had a good acquisition history, good crop calendar information, and was one of the first segments to pass through pre-processing (sun angle and haze correction, etc.). GRABS values were used to calculate the detectability date because GRABS=0 (green vegetation threshold value) is standard for all segments and acquisitions processed through the LACIE Program. Mean GRABS values of ten soybean fields were plotted for the first five acquisitions: June 1, June 10, July 16, July 26 and August 4. All ten fields passed the GRABS threshold of detectability between the fourth and fifth acquisitions, Figure 5. A variable x (Julian date) = the t intercept, of the line drawn between GRABS values on acquisitions 4 and 5 with the threshold of detectability (GRABS = 0), was determined for each sampled field. Using the value of this variable for each sample field, a weighted mean and variance was calculated. The 95% confidence interval was determined and added to the weighted mean giving a Julian day by which 95% of sampled soybean fields were detectable. In 1978 50% statewide planted day for soybean was obtained from the Indiana annual crop and livestock report for the year and subtracted from the 95% mean detectability day + the 95% confidence interval. The difference, 15 days, was the interval x_2 in Figure 4.

The interval from the 50%-planted day for corn and the 50%-planted day for soybean was ascertained as shown in Table 4. Six years of crop calendar data was available from which to calculate a mean number of days from the statewide 50% corn planted day to the 50% soybean-planted day. The 95% confidence interval was determined and added to the mean to give an interval, x_1 in Figure 4, of 26 days. The 26-day interval was added to the 15-day interval from soybean planting to first detectability to estimate x_1 in Figure 4--the number of days from the 50%-planted day for corn to first detectability of all soybean fields in segment 127. The final value, 41 days, was rounded off to 40 days for convenience.

Although the 40-day interval from corn planting to soybean detectability was calculated from a single segment year, the time interval $t_0 + 40$ was used as the initial estimate of $(t_2 - t_0)$ for all 18 segments. The actual value for $t_2 - t_0$ varied from state to state because of state-to-state variation in the 50% corn-planted day, but almost all corn and soybean fields were detectable on the first acquisition after $t_0 + 40$.

99.1% of summer crop fields had mean GRABS values above 0 on the first acquisition after t_0 . 97.3% of the fields had mean values above GRABS = 5.0. The number of days from t_0+40 to the first acquisition varied between 5 and 40 with an average, weighted by the number of fields in each segment, of 18.5 days. In contrast, 41.8% of summer crop fields had GRABS values above 0 on the last acquisition prior to the t_0+40 date. The number of days between the acquisition prior to t_0+40 and the (t_0+40) date varied between 5 and 31 with a weighted average of 15.0. The data is summarized in Table 5.

The estimate of a 40-day interval from planting to detectability was further evaluated using field measurements data obtained from LARS. Corn fields had at least 20% ground cover* approximately 40 days after planting, especially with the more dense, and more common, plantings. Soybean fields passed the 20% ground cover point well before 40 days after 50% planted date of the soybeans. The spectral emergence of the soybean fields could not be evaluated with respect to a 50% corn planted date since this was somewhat of an artificial situation.

Additional analysis was performed to estimate t_2 using all corn and soybean pixels in seven Indiana and seven Iowa segments. Digitized and registered ground data was used to sample all identified corn and soybean pixels within the 14 segments. The percentage of corn and soybean pixels above GRABS = 5.0 decision threshold** on each acquisition for each segment was determined and plotted against Julian date. The segment plots were then combined by state and a statewide (as represented by the sample of seven segments/state) cumulative spectral emergence curve (S curve) was generated for each state. The S curve was fit to the data points using a two piece curve fitting method whereby all data before the obvious upper break in slope plus the 1st group of data points at or after the obvious break in slope were fitted by regression to the exponential function $y = ae^{bx}$. The second group of points which included all data points at or after the obvious upper break in slope were fit by regression to the linear function $y = ax+b$. The two curves were then pieced together visually (i.e. the upper shoulder was estimated by eye). Two analyses were done, a) one using t_0 = 50% planted date for corn statewide, and b) one using t_0 = 50% planted date for corn at the CRD to which a given segment belonged. Thus the second analysis was done on data normalized for t_0 at the CRD level. The non-normalized S curves of data points for each set of segments by state are shown in Figure 6. The normalized S curves and data points are shown in Figure 7.

The results of this analysis are shown in Table 6. The earlier estimate of $t_0+40 = t_2$ = 95% summer crops spectral emerged based on analysis of segment 127 sampled corn and soybean fields does not appear to have been very good. From the above described analysis $t_2 = t_0+70$ and t_0+51 for Iowa and Indiana non-normalized data respectively. For the normalized

* From work at ERIM it has been estimated that approximately 20% ground cover is necessary in order to reach the threshold of spectral detectability.

** Two GRABS detectability thresholds were used, 0 and 5, because temporal plots of summer crop field mean values indicated that GRABS values sometimes fluctuated between 0 and 5 prior to t_0+40 . GRABS values did not exceed 5 on any consecutive acquisitions prior to t_0+40 .

data, $t_2 = T_0 + 58$ and $T_0 + 56$ days for Iowa and Indiana respectively. There was also better correspondence between estimates of other % emergence points between the two states in the normalized data. The r^2 values for the regression curve fit were fairly good for Indiana - .86 and .84 for non-normalized and normalized data respectively. However the r^2 values for the regression curve fit for Iowa were not very good - .57 and .46 non-normalized and normalized data respectively.

This new estimate for $t_2 = t_0 + 57$ days actually corresponds very well with the definition for the end of Summer Crop biowindow SC2 from C/S-1 baseline procedure if one adjusts for the difference between 5% corn planted (from C/S-1) and 50% corn planted from the Through-the-Season Model. This was to be expected since the SC2/SC3 biowindow boundary point and t_2 are theoretically the same from their conceptual definitions.

It is clear from inspection of Figures 6 and 7 that there is much more variation in Iowa spectral crop calendars. To what this variation is due cannot be determined in any certainty in this study, however it would seem reasonable to suspect that most of the crop calendar variation in Iowa is due to greater variation in planting dates throughout the state than occurs in Indiana. The observation that the two normalized fitted spectral emergence curves for Iowa and Indiana were similar leads one to suspect that normalization for planting date variation can be valuable for any agricultural multitemporal spectral data processing. This of course was the idea behind UCB's dynamic stratification work. However the large variation in the Iowa data even after normalization seems to suggest that crop calendar "strata" in Iowa are smaller than CRD size. In Indiana on the other hand CRD size strata seem to be reasonable.

The above observations are relevant to the Through-the-Season effort in that it suggests that any Through-the-Season Labeling Procedure or Area Estimation Procedure based on certain kinds of models (i.e. models that have some variable based on crop calendar) would be more effective if normalization for planting date can first be performed. Another method for controlling the variation would be through crop calendar stratification.

Which method would be more desirable would be totally dependent upon the over-all area estimation procedure design and the operational environment of the procedure.

6.0 EARLY SEASON TEMPORAL-SPECTRAL CHARACTERISTICS: SUMMER CROPS VS OTHER

Edwin J. Sheffner

6.1 GREEN VEGETATION INDICATOR-GRABS

Differences between summer crops and non-summer crop temporal profiles become more apparent when examined within the context of the Through-the-Season Labeling Accuracy Model where t_2 is defined as a reference point. Note: The following analysis was performed prior to the latest estimation of $t_2 = t_0 + 57$ days, and so the analysis was performed relative to the original estimate of $t_2 = t_0 + 40$ days.

6.1.1 Summer Crops vs. Other

Spectral Emergence

Mean GRABS values for corn, soybean and sorghum fields rarely exceeded the decision threshold GRABS = 5.0 on acquisitions prior to the t_0+40 date. 1572 was the only segment with summer crop mean GRABS values above 5.0 on any acquisition prior to t_0+40 (sorghum), and no segments had two consecutive acquisitions prior to t_0+40 with summer crop values above 5.0. Sixteen of the 18 sampled segments had alfalfa/pasture fields. Mean GRABS values of alfalfa/pasture exceeded 5.0 on at least one acquisition prior to t_0+40 in all 16 segments. Fourteen of the 16 segments with alfalfa/pasture fields had more than two acquisitions prior to t_0+40 ; in 12 of the segments alfalfa/pasture field mean GRABS values exceeded 5.0 on at least three acquisitions prior to t_0+40 .

Idle cover and fallow fields also tended to have higher GRABS values than summer crops early in the season. Eight segments had sampled idle cover fields and six segments had sampled fallow fields. Idle cover field mean GRABS values exceeded 5.0 on more than one acquisition prior to t_0+40 in five of the eight segments and exceeded 5.0 on more than two acquisitions prior to t_0+40 in four of the eight segments. Fallow field mean GRABS values exceeded 5.0 on more than two acquisitions prior to t_0+40 in three of six segments.

The sampled summer crop confusers tended to reach maximum GRABS values earlier in the season than summer crops. All summer crop fields reached maximum GRABS values after the t_0+40 date, but alfalfa/pasture fields reached peak GRABS values prior to t_0+40 in 12 of 16 segments, idle cover fields reached peak GRABS values prior to t_0+40 in two of six segments.

A dip in summer crop mean GRABS values prior to the t_0+40 date occurred in several segments. The GRABS temporal plots of corn and soybean in segment 127, (Figure 5), illustrate the dip in GRABS values which occurred between the third and fifth acquisitions. The GRABS value dip was a more common occurrence in soybean and sorghum fields than in corn fields. The dip occurred in soybean fields in 14 of 17 segments and in three of four segments with sorghum, but only in half the segments (9 of 18) with corn. Similar dip phenomenon was noted in non-summer crops. Nine of 16 segments with alfalfa/pasture, two of eight segments with idle cover fields and two of six segments with fallow fields had similar dips prior to t_0+40 . The cause of the dip is uncertain. It may be a response to cultivation practices or weather or variation in sensor calibration. In either case, its appearance in the spectral data is acquisition dependent.

6.1.2 Differences Between Summer Crops

The GRABS temporal plots showed differences among the summer crops in timing and range of peak GRABS values. Corn fields generally peaked earlier than soybean fields and sorghum fields were mixed. Mean corn

field values reached a peak between t_0+40 and t_0+85 in 11 of 17 segments, whereas soybean fields reached peak GRABS values after date t_0+85 in 10 of 15 segments. Three segments lacked adequate acquisitions to use in this analysis. Segments with sorghum fields were evenly split--two of the four segments had sorghum fields which peaked in the period t_0+40 and t_0+85 and two segments had sorghum fields which peaked in the period after t_0+85 . During specified periods of the growing season, mean summer crop GRABS differed. In the period between t_0+40 and t_0+57 , corn mean GRABS values varied between 14.42 and 31.12. Mean soybean field values were lower, varying between 1.03 and 27.62. Later in the season, the situation reversed. For acquisitions obtained between t_0+85 and t_0+102 , mean corn values were between 17.62 and 22.63. Soybean mean field values were between 31.52 and 45.65 during the same period. Sorghum field GRABS values varied between 7.17 and 12.18 in the earlier period and between 15.67 and 31.12 later in the season.

GRABS emerged as a significant feature for early season separation of summer and non-summer crops. Vegetation within fields of major summer crop confusers tend to be detectable on Landsat in the GRABS feature earlier than summer crops, reach peak values before summer crops, and generally described different spectral development patterns. The descriptions of the development of the various cover types given above and the conclusions drawn from them are based on mean values of all field means of a given cover type. Individual fields varied significantly from the mean as can be seen in Appendix A by the plots of standard deviation. Some non-summer crop fields, particularly idle cover and fallow fields, mimic summer crops in GRABS development pattern, and, occasionally, a field identified as summer crop on ground data was observed to follow a typical alfalfa/pasture pattern. The possibility that anomalies are present must be kept in mind during the labeling process.

6.2 BRIGHTNESS VALUES

The dip noted in the GRABS feature for summer crops is also observed in the tassell-cap brightness feature. Mean TC1 values for corn fields in 12 of 17 segments dipped and then increased in value prior to t_0+40 . The same behavior occurred for soybean fields in 14 of 16 segments and 3 of 4 segments with sorghum. The TC1 dip also occurred in some non-summer crop fields. Mean values for idle cover fields in five of eight segments and for fallow fields in 4 of 6 segments had a TC1 dip, but the dip was evident in only 5 of 15 segments with alfalfa/pasture.

The time of maximum TC1 values varied among the summer crops and between the summer crops and non-summer crops. Maximum TC1 values occurred prior to t_0+40 in corn fields in 13 of 17 segments and in 3 of 4 segments with sorghum fields, but only in 4 of 16 segments with soybean. Maximum soybean TC1 values were attained between t_0+40 and t_0+85 in 6 segments and after t_0+85 in 6 additional segments.

Maximum TCI values were reached in half the segments with alfalfa/pasture fields prior to t_0+40 and between t_0+40 and t_0+85 in the majority of the remaining alfalfa/pasture segments.

TCI values for soybeans remained consistently higher than corn or sorghum values after the t_0+40 date. Soybean field mean values of TCI varied between 56.38 and 81.68 on acquisitions between t_0+40 and t_0+57 and between 75.10 and 90.74 for acquisitions between t_0+85 and t_0+102 . Corn field values for the same periods varied between 61.73 and 74.98, and 61.79 and 83.76 respectively. Sorghum TCI values were similar to corn in both time frames.

7.0 SUMMARY

Landsat data has proven to be effective in large scale agricultural inventories. However, prior to AgRISTARS, the usefulness of Landsat data was limited by the lack of labeling logic for identification of crops other than small grains and earlier than at-harvest.

To improve the timeliness of Landsat area estimates, UCB undertook the experiment described in this report. The results of the experiment included a framework for understanding how Landsat data can be used in conjunction with ancillary data to identify crop groups (in the Corn Belt) at any time during the growing season (labeling accuracy plateaus) and the demonstration that first detectability of green vegetation on Landsat was a critical event in the spectral development of summer crops in the Corn Belt which could be used to separate summer crops from non-summer crops as early as 40 days after the 50%-planted date statewide for corn (t_0+40). Maximum green development was identified as the earliest time in the growing season when corn and soybeans are separable.

While the results of this experiment indicate that procedures for summer crop labeling prior to harvest are feasible, there are a number of significant problems that remain to be examined. The method described in this report for determining the boundaries of the labeling accuracy plateaus assume that a season-specific starting date (from crop calendar information or a planting model) will be available. The assumption may not be realistic.

Although t_0+40 may have no meaning outside the Corn Belt, the labeling accuracy plateau approach may be appropriate in other regions. The nature of summer crop cultivation in non-Corn Belt regions needs to be studied before Through-the-Season labeling procedures can be written.

Within the Corn Belt, additional research is required to quantitatively establish the remaining boundaries of the labeling accuracy plateaus. A multiyear approach which incorporates information on crop rotation and planting patterns may yield a reliable date prior to t_0+40 for summer crop/non-summer crop discrimination.

TABLE 1. List of "Important Questions" for Crop Type Identification

Question #	Question Statement	Determinant Characteristic
1	When (during what time period) is a green vegetation canopy detectable or present within the given feature?	Temporal
OR THREE MORE SIMPLE, ALTERNATIVE QUESTIONS:		
1a	Is there a bare soil stage observable? When?	Temporal
1b	When is a green vegetation canopy first detectable within the spectral data for a given feature?	
1c	When is a green vegetation canopy last detectable within the spectral data for a given feature?	
2	When do the maximum (or minimum) values for specific spectral variables occur?	Temporal
2a	When does the maximum value for the measure of "infraredness," the green vegetation indicator, occur?	Temporal
2b	When does the maximum or minimum value for the brightness measure (Brightness) occur?	Temporal
2c	When does the maximum or minimum value for the visual yellow-green measure occur? (Can not be adequately measured with Landsat at this time)	Temporal
3	What is the value of the maximum for the Green Vegetation Indicator?	Spectral
4	Is there a senescent biostage observable for the feature?	Spectral
5	If there is a senescent biostage observable for the feature, when does it occur?	Temporal
6	What is the shape of the temporal profile (Temporal plot) of the green vegetation indicator vs. time? (Does the profile have a plateau, is it broad-peaked, narrow-peaked, tall, squat, etc.?)	Temporal-Spectral

TABLE 2. Location and Acquisition Dates - Data Set Segments

Segment	Location	Acquisition Dates								
123	Hamilton Co. Indiana	89	107	152	161	197	233	269		
127	Montgomery Co. Indiana	89 269	107	152	161	197	207	216	243	252
141	Madison Co. Iowa	86	103	130	166	167	212	220		
144	Wapello Co. Iowa	130	165	183	219	220	238	246	264	274
205	Clark Co. Missouri	93 282	101 290	137	155	164	209	218	246	272
809	Ogle Co. Illinois	101	163	164	209	218	244	254		
832	Adams Co. Indiana	88	97	151	160	178	232	268		
843	Henry Co. Indiana	88	97	151	152	160	178	197	251	
852	Randolf Co. Indiana	88	97	151	160	178	232	250	268	
853	Randolf Co. Indiana	88	97	151	160	178	232	250	268	
860	Wells Co. Indiana	88 197	97 232	107 251	116	151	152	160	161	178
864	Crawford Co. Iowa	87 258	96	141	150	159	186	222	231	248
865	Crawford Co. Iowa	87 267	96	141	153	159	168	186	231	249
880	Monona Co. Iowa	87	96	141	150	186	204	222		
881	Monona Co. Iowa	87 267	96	141	159	186	213	222	231	249
882	Palo Alto Co. Iowa	86 231	96 258	131 267	141 293	150	159	186	213	221
1572	Custer Co. Nebraska	134	153	170	189	206	224	243		
1591	Webster Co. Nebraska	134	152	169	188	205	223	241	259	

TABLE 3: Number of Fields Sampled From Data Set

SEGMENT	CROP TYPE					CROP TYPE					
	C	SY	SR	SB	SU	A	P	H	IC	F	T
123	7	5				3			3	3	1
127	10	10			2	4		2	3	1	3
141	7	6				4	5	4			
144	10	10				3	5	5			5
205	10	9		2			1	1	1	6	4
809	10	10				5	3	3			
832	10	10				5	2	3		2	
843	9	10				3	5	2	2		
852	10	10				2	5	4		2	
653	10	10				2	3	3			
860	8	8				1		2		2	
864	6	10				4	6	2	3		
865	10	10	5				7		2		
880	10	10	4			2	2		15	3	
881	10	10				7	5		9	1	5
882	10	10				2	8		3	2	
1572	5	0	5			5	5			5	
1591	10	3	7				5		3	1	
TOTAL	162	151	21	2	2	52	67	31	34	28	17

TOTAL: Summer Crops = 338 Non-Summer Crops = 236

C = Corn	A = Alfalfa
SY = Soybean	P = Pasture
SR = Sorghum	H = Hay
SB = Sugar Beet	IC = Idle Cover
SU = Sunflower	F = Fallow
	T = Trees

Table 4: Calculation of maximum number of days between 50% planted day of Corn and 50% planted day of soybean in Indiana.

Year	A Day on which 50% of Corn was Planted*	B Day on which 50% of Soy was Planted*	B-A
1973	140	161	21 days
1975	133	140	7
1976	124	140	16
1977	131	142	11
1978	152	160	8
1979	138	148	10
	<u>136.33 ± 24.48</u>	<u>148.5 ± 25.07</u>	

Mean = 12.17 days

Standard Deviation = 5.34 days

NOTE: 95% Confidence Interval = 12.17 days ± 13.74

Day Interval = 12.17 + 13.74 = 25.91 or 26 days

* From Indiana Annual
Crop and Livestock
Summaries

TABLE 5: Accuracy to $t_0 + 40$ Date in Predicting Summer Crop Detectability on Landsat

Segment	$t_0 + 40$ Day*	A	B	C	D
123	192	5	12	12 (100)	12 (100)
127	192	5	20	20 (100)	20 (100)
141	174	38	13	13 (100)	13 (100)
144	174	9	15	15 (100)	15 (100)
205	178	31	21	21 (100)	21 (100)
809	184	25	20	20 (100)	20 (100)
832	192	40	20	20 (100)	20 (100)
843	192	5	20	20 (100)	19 (95.0)
852	192	40	20	20 (100)	20 (100)
853	192	40	20	20 (100)	20 (100)
860	192	5	16	16 (100)	15 (93.8)
864	174	12	16	16 (100)	16 (100)
865	174	12	24	24 (100)	24 (100)
880	174	12	23	23 (100)	23 (100)
881	174	12	20	19 (95.0)	18 (90.0)
882	174	12	20	20 (100)	20 (100)
1572	175	14	10	8 (80.0)	8 (80.0)
1591	175	13	20	20 (100)	20 (100)

A = Number of days from $t_0 + 40$ date to first acquisition after $t_0 + 40$.

B = Number of sampled summer crop fields.

C = Number of sampled summer crop fields with mean GRABS values above GRABS=0 on first acquisition after $t_0 + 40$ (%).

D = Number of sampled summer crop fields with mean GRABS values above GRABS=5 on first acquisition after $t_0 + 40$ (%).

* = $t_0 + 40$ Day = 40 days after 50% corn planted day, statewide.

Summary of Above Data

Number of Segments

4	5	68	68 (100)	66 (97.1)
1	9	15	15 (100)	13 (86.7)
5	12	103	102 (99.0)	101 (98.1)
2	13/14	30	28 (93.3)	27 (90.0)
6	20+	114	114 (100)	114 (100)

TABLE 6: Evaluation of $(t_2 - t_0)$ using accumulated 's-curve' approach.

% Corn + Soybean Spectrally Emerged	$t_{\% \text{ emerged}} - t_0$ (days)		$t_{\% \text{ emerged}} - t_0$ (days)	
	Non-Normalized t_0 = statewide 50% planted date of corn		Normalized t_0 = CRD 50% planted date of corn	
	IOWA	INDIANA	IOWA	INDIANA
95%	70 days	51 days	58 days	56 days
90%	57 days	45 days	52 days	52 days
75%	49 days	41 days	47 days	48 days
50%	42 days	33 days	39 days	40 days
25%	29 days	17 days	25 days	25 days

Coefficients from $y = ae^{bx}$ curve fit

a	6.95	10.38	7.22	7.91
b	.05	.05	.05	.05
r^2	.57	.86	.46	.84




FIGURE 1. General, through-the-season, labeling accuracy model.




FIGURE 2. Specific, through-the-season, labeling accuracy model for corn and soybeans in the U.S. Corn Belt and peripheries.

ORIGINAL PAGE IS
OF POOR QUALITY

FIGURE 3

LOCATION OF SAMPLE SEGMENTS





Figure 4

Method for Determining when Summer Crops Have Passed
Landsat Threshold of Green Vegetation Detectability

50% PLANTING DATE FOR CORN*	DATE BY WHICH 50% OF SOY PLANTED IN 95 OUT OF 100 YEARS**	DATE BY WHICH 95% OF SOY FIELDS DETECTABLE
--------------------------------	---	--




ΔT = NUMBER OF DAYS AFTER 50% PLANTING DATE
FOR CORN WHEN WINDOW FOR EARLY SEASON
ESTIMATE CLOSES

$$\Delta T = x_1 + x_2$$

*FROM CROP CALENDAR
OR PLANTING MODEL

**FROM CROP CALENDAR DATA



Acquisitions

MR30 (89)
AP17 (107)
JN01 (152)
JN10 (161)
JY16 (197)
JY26 (207)
AG04 (216)

Figure 5

GRABS/Temporal Plots of
Corn and Soybean Field Means, Segment 127

X-axis scales are equal. All fields are detectable
40 days after 50% planted day for corn.

FIGURE 6. Percentage of Corn and Soybean Pixels above GRABS = 5.0 as a Function of Time from Statewide Corn 50% Planted Date.







FIGURE 7. Percentage of Corn and Soybean Pixels above GRABS = 5.0 as a Function of Time from Statewide Corn 50% Planted Dated Adjusted to CRD Level.



REFERENCES

1. Cicone, R., C. Hay, R. Horvath, M. Metzler, O. Mykolenko, J. Odenweller, and D. Rice, 1981. Users Manual for the U. S. Baseline Corn and Soybean Segment Classification Procedure. NASA Technical Report #FY-EI-00712, Environmental Research Institute of Michigan and University of California, Berkeley.
2. Kauth, R. J., P. F. Lambeck, W. Richardson, G. S. Thomas, and A. P. Pentland, 1979. "Feature Extraction Applied to Agricultural Crops as Seen by Landsat," in The LACIE Symposium. JSC-16015, NASA/Johnson Space Center, Houston, Texas.
3. Hay, C., C. Kuretz, J. Odenweller, E. Sheffner, and B. Wood, 1979. Development of AI Procedures for Dealing with the Effects of Episodal Events on Crop Temporal Spectral Response and Development of AI Guidelines for Corn and Soybean Labeling. Final Report for NASA Contract NAS9-14565, SSL Series 20, Issue 44, Berkeley, California.

APPENDIX A


GRABS and Brightness versus Time Plots

Plots of GRABS (A-1 to A-36) and Brightness (A-37 to A-72) versus time for corn, soybeans and other crops for sample segments listed in Table 2. The actual values equal the plotted value minus 10. The horizontal line indicates the detection threshold (GRABS=0). The first vertical line equals $t_0 + 40$; the second vertical line equals $t_0 + 85$.


ORIGINAL PAGE IS
OF POOR QUALITY

A-1

127 CORN
GRAB
-1 STANDARD DEVIATION. MEAN. ± 1 STANDARD DEVIATION



123 CORN
GRAB
-1 STANDARD DEVIATION. MEAN. ± 1 STANDARD DEVIATION




ORIGINAL PAGE IS
OF POOR QUALITY

A-2

ALL COM
GRAB
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

75.00000
63.30320
51.60656
39.90304
28.20311
16.50639




ALL COM
GRAB
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

0.0 37 117 157 197 237


0.0 37 117 157 197 237

201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300

NOIADJ30 OVRSTATS 1- NEW NOIADJ30 OVRSTATS 1-
58925
MM03 582



MOBILE 0204515 T, WASH, 01 STANFORD UNIVERSITY 1-
5085
MAY 68



2


[illegible]

64.....65.....66.....67.....


61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 10

ORIGINAL PAGE IS
OF POOR QUALITY

9-3 COM
2825
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



9-2 COM
2825
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



79.00300

53.30129

31.60551

99.9874

29.21311

16.51539


0.0

50009
MAR 1966

MEAN, ± 1 STANDARD DEVIATION


NOIIVIA3G OTTCNVIS 10 'WCM 'NCIIVIA3G OTTCNVIS 10-
568

ORIGINAL PAGE IS
OF POOR QUALITY



A-5


1-1 STANDARD DEVIATION. MEAN. +1 STANDARD DEVIATION



41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871


ACQUIRED IMMUNITY TO TUBERCULOSIS DUE TO

A-6

[illegible]

865 CORN
GRAB
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION


75.0000
63.30320
51.60656
39.90996
28.21311
16.51639



0.0

890 CORN
GRAB
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

75.0000
63.30320
51.60656
39.90996
28.21311
16.51639



0.0

A-7

ORIGINAL PAGE IS
OF POOR QUALITY


21.774
Ym

982 CORN
GRAB
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

ORIGINAL PAGE IS
OF POOR QUALITY

A-8


75.00000
63.30320
51.60656
39.90992
28.21311
16.51639



0.0
3.....37.....77.....117.....157.....197.....237

481 CORN
GRAB
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION


75.00000
63.30320
51.60656
39.90992
28.21311
16.51639




0.0
3.....37.....77.....117.....157.....197.....237

1972 CORN
CR805
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

ORIGINAL PAGE IS
OF POOR QUALITY




1972 CORN
CR805
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



ORIGINAL PAGE IS
OF POOR QUALITY

127 SOYBEANS
64695
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION


75.00000
65.30320
51.00556
39.90904
20.21311
16.51639



0.0 10 20 30 40


123 SOYBEANS
63005
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

75.00000
65.30320
51.00556
39.90904
20.21311
16.51639




0.0 10 20 30 40

144 INCHES
CROSS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




144 INCHES
CROSS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



ORIGINAL PAGE IS
OF POOR QUALITY


205 5076243
22485
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



512 5075 845

22-35

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




A-13

ORIGINAL PAGE IS
OF POOR QUALITY

512 5075 845


22-35

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




022 SOYBEANS
CRASS

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



023 SOYBEANS
CRASS

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



21 2021
VIRGINIA 10


ORIGINAL PAGE IS
OF POOR QUALITY

ORIGINAL PAGE IS
OF POOR QUALITY

A-15


833 SOYBEANS
2285

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



833 SOYBEANS
2285


-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




0.0 37 77 117 157 197

0.0 37 77 117 157 197

ACTIVATION CORRELATES TO STANDARD DEVIATION. MEAN. 1 STANDARD DEVIATION. 1-2



SEVEN
SNV30105 000

-1- STANDARD DEVIATION. MEAN. ± 1 STANDARD DEVIATION


ORIGINAL PAGE IS
OF POOR QUALITY

A-18

1991 SURVEILANCE
CRABS

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

75.00000
63.30320
51.60636
39.90956
28.21316
16.51639




0.0

45
44
43
42
41
40
39
38
37
36
35
34
33
32
31
30
29
28
27
26
25
24
23
22
21
20
19
18
17
16
15
14
13
12
11
10
9
8
7
6
5
4
3
2
1
0

1992 SURVEILANCE
CRABS

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

75.00000
63.30320
51.60636
39.90956
28.21316
16.51639



0.0

45
44
43
42
41
40
39
38
37
36
35
34
33
32
31
30
29
28
27
26
25
24
23
22
21
20
19
18
17
16
15
14
13
12
11
10
9
8
7
6
5
4
3
2
1
0

ORIGINAL PAGE IS
OF POOR QUALITY

A-19

127 MAY/PASTURE
GRASS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

75.00000

63.30320

51.60656

39.90992

28.21311

16.51639

0.0

-3

37

77

117

157

197

237

0.0
-3
37
77
117
157
197
237

123 ALFALFA/MAY/PASTURE
GRASS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

75.00000

63.30320

51.60656

39.90992

28.21311

16.51639

0.0

-3

37

77

117

157

197

237

25.00000	63.39320	51.60455	39.99001	20.21111	16.91639
----------	----------	----------	----------	----------	----------





Figure 1

3

75.00000	63.30320	91.48656	39.90904	20.21311	16.51619
----------	----------	----------	----------	----------	----------




453
.....451
.....451
.....411
.....44
.....45
.....4


3

ORIGINAL PAGE IS
OF POOR QUALITY

032 ALPHE/PMV/PASTURE
C8835
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



032 ALPHE/PMV/PASTURE
C8835
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



032 ALFALFA/HAY/PASTURE
GRASS

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

031 ALFALFA/HAY/PASTURE
GRASS

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

ORIGINAL PAGE IS
OF POOR QUALITY

75.00000



63.30320

51.60656

39.90992

20.21311

10.51539



0.0

3.....11.....117.....157.....197.....237.....



-1.....19.....35.....51.....67.....83.....99.....115.....131.....147.....163.....179.....195.....

MAIL ROOM 50423
2000/04/09 14:14 896

1-STANDARD DEVIATION, MEAN, \pm 1 STANDARD DEVIATION

ORIGINAL PAGE IS
OF POOR QUALITY

A-23




ORIGINAL PAGE IS
OF POOR QUALITY

855 PASTURE
GRASS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

75.0000
63.39320
51.68556
39.39906
26.21311
16.51039


A-24



0.0
10
20
30
40
50
60
70
80
90
100


136 ALFALFA/PASTURE/GRASS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

75.0000
63.39320
51.68556
39.39906
26.21311
16.51039



0.0
10
20
30
40
50
60
70
80
90
100


880 ALFALFA/PASTURE
CRABS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



0.0 37 117 137 157 237

881 ALFALFA/PASTURE/CROSS
CRABS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION


ORIGINAL PAGE IS
OF POOR QUALITY



0.0 37 117 137 157 237

052 GRASS/HAY/PASTURE
GRASS

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




0.0 3 37 77 117 157 197 237

1972 ALVALFA/PASTURE
GRASS

ORIGINAL PAGE IS
OF POOR QUALITY

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




0.0 3 37 77 117 157 197 237

A-27

332 FALLOU FIELD
GRASS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

75.0000
63.38320
51.0656
39.9090
20.21311
16.51639




332 FALLOU FIELD
GRASS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

0.0

123 FALLOU
GRASS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

75.0000
63.38320
51.0656
39.9090
20.21311
16.51639




0.0

ORIGINAL PAGE IS
OF POOR QUALITY

000 FALLON
CROSS

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION


79.00000
63.30320
51.40696
39.90906
20.21311
10.91639



000 FALLON
CROSS

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

79.00000
63.30320
51.40696
39.90906
20.21311
10.91639



0.0


0.0

ORIGINAL PAGE IS
OF POOR QUALITY

1972 FALLON
CRAB
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION


75.00000
63.30320
51.60656
39.90992
28.21311
16.51539

A-29



1982 FALLON
CRAB
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

75.00000
63.30320
51.60656
39.90992
28.21311
16.51539



0.0
117
157
197

ORIGINAL PAGE IS
OF POOR QUALITY



A-30

123 ILE COVER
CROSS

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

123 ILE COVER
CROSS


-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




ORIGINAL PAGE IS
OF POOR QUALITY

8-1 IDLE COVER GRP
GRAB
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

A-31



8-1 IDLE COVER
GRAB
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




0.0
3.0
10.0
15.0
20.0
25.0
30.0
35.0
40.0
45.0
50.0
55.0
60.0
65.0
70.0
75.0
80.0
85.0
90.0
95.0
100.0

75.00000
63.10320
51.60656
39.93906
29.21311
16.91639

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



MEAN, ± 1 STANDARD DEVIATION, ± 1 STANDARD DEVIATION





891 IDLE COVER
G2105

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

892 IDLE COVER
G2105

-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




A-33

ORIGINAL PAGE IS
OF POOR QUALITY


ORIGINAL PAGE IS
OF POOR QUALITY

127 SUNFLOWERS
STUBS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




ORIGINAL PAGE IS
OF POOR QUALITY

888 508CHUM
CR485
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION





A-35




8.0 117 157 197 237

883 508CHUM
CR485
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



8.0 117 157 197 237

1572 SUGCMM
 CRAB
 -1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



0.0 0.5


1591 SUGCMM
 CRAB
 -1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

ORIGINAL PAGE IS
 OF POOR QUALITY



0.0

A-36




0.0 0.5


ORIGINAL PAGE IS
OF POOR QUALITY

A-37

127 COM
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION





125 COM
41
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



NOTED BY BUREAU TO ADV. NO. 10-10-58
SEEKING TO
NO. 582

164 CORN
KI-BRIGHTNESS
STANDARD DEVIATION: MEAN: +1 STANDARD DEVIATION



A-38



.....

.....46
.....47
.....48
.....49
.....50


ORIGINAL PAGE IS
OF POOR QUALITY

A-39

022 02M
CL-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



022 02M
CL-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




0.0 1.0 2.0 3.0 4.0 5.0 6.0 7.0 8.0 9.0 10.0

0.0 1.0 2.0 3.0 4.0 5.0 6.0 7.0 8.0 9.0 10.0


022 02M
CL-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

ORIGINAL PAGE IS
OF POOR QUALITY

833 200W
K1-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



932 CORN OIL
K1-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




0.0 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

ORIGINAL PAGE IS
OF POOR QUALITY

000 200M
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

125.3000
105.1066
85.21311
65.31967
45.42623
25.93279




0.0

OF PHOTO COPY

001 200M
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

125.3000
105.1066
85.21311
65.31967
45.42623
25.93279




0.0


ORIGINAL PAGE IS
OF POOR QUALITY

A-42

845 5004
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



845 5004
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



0.0

0.0

ORIGINAL PAGE IS
OF POOR QUALITY


A-43

001 CORN
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



0.0 37 77 117 157 197 237

001 CORN
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




0.0 37 77 117 157 197 237


ORIGINAL PAGE IS
OF POOR QUALITY

A-44

1972 CORN
CI-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



1982 CORN
CI-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



0.0 1.0 2.0 3.0 4.0 5.0 6.0 7.0 8.0 9.0 10.0


0.0 1.0 2.0 3.0 4.0 5.0 6.0 7.0 8.0 9.0 10.0

123 SOYBEANS

41

-1 STANDARD DEVIATION. MEAN, +1 STANDARD DEVIATION


125.0000
105.1066
85.2131
65.3197
45.4262
25.5327



0.0

.....45.....47.....49.....51.....53.....55.....57.....59.....61.....63.....65.....67.....69.....71.....73.....75.....77.....79.....81.....83.....85.....87.....89.....91.....93.....95.....97.....99.....100

125.0000
105.1066
85.2131
65.3197
45.4262
25.5327



0.0

.....45.....47.....49.....51.....53.....55.....57.....59.....61.....63.....65.....67.....69.....71.....73.....75.....77.....79.....81.....83.....85.....87.....89.....91.....93.....95.....97.....99.....100

ORIGINAL PAGE IS
OF POOR QUALITY

A-45



127 SOYBEANS
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

ORIGINAL PAGE IS
OF POOR QUALITY

A-46

208 507 BEAMS
41-2K15H15-SS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

171 307E-11
41-2K15H15-SS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



125.0000

103.1056

85.21311

65.31367

45.62623

25.93279

0.0

37 77 117 157 197 237

ORIGINAL PAGE IS
OF POOR QUALITY

A-47

002 SURFACES
CL-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

.....

125.0000


105.1062

85.21311

65.31967

-5.42623

25.53279



.....

0.0

003 SURFACES
CL-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

.....

125.0000


105.1066

85.21311

65.31967

-5.42623

25.53279




.....

0.0


ORIGINAL PAGE IS
OF POOR QUALITY

A-48

922 507MEANS
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



922 507MEANS
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



125.0000

105.1066

05.21311

05.31967


49.42623

29.53279

0.0

033 20424MS
K1-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION


125.0000
105.1565
85.21311
65.31967
45.62623
25.93279



0.137.....77.....117.....157.....197.....237.....

068 20424MS
K1-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

125.0000
105.1065
85.21311
65.31967
45.62623
25.93279



0.037.....77.....117.....157.....197.....237.....


ORIGINAL PAGE IS
OF POOR QUALITY

A-49


ORIGINAL PAGE IS
OF POOR QUALITY

A-50

85. SOURCE
AL-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




85. SOURCE
AL-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



ORIGINAL PAGE IS
OF POOR QUALITY


A-51

031 503445
K1-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




0.0 0.1 0.2 0.3 0.4 0.5 0.6 0.7 0.8 0.9 1.0

030 503445
K1-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




0.0 0.1 0.2 0.3 0.4 0.5 0.6 0.7 0.8 0.9 1.0

1941 SYMBOLS
CL-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



1942 SYMBOLS
CL-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION





37 77 117 157 197 237

37 77 117 157 197 237

886 50462855
KI-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

ORIGINAL PAGE IS
OF POOR QUALITY

A-53.




0.0
1.0
2.0
3.0
4.0
5.0
6.0
7.0
8.0
9.0
10.0
11.0
12.0
13.0
14.0
15.0
16.0
17.0
18.0
19.0
20.0
21.0
22.0
23.0
24.0
25.0
26.0
27.0
28.0
29.0
30.0
31.0
32.0
33.0
34.0
35.0
36.0
37.0
38.0
39.0
40.0
41.0
42.0
43.0
44.0
45.0
46.0
47.0
48.0
49.0
50.0
51.0
52.0
53.0
54.0
55.0
56.0
57.0
58.0
59.0
60.0
61.0
62.0
63.0
64.0
65.0
66.0
67.0
68.0
69.0
70.0
71.0
72.0
73.0
74.0
75.0
76.0
77.0
78.0
79.0
80.0
81.0
82.0
83.0
84.0
85.0
86.0
87.0
88.0
89.0
90.0
91.0
92.0
93.0
94.0
95.0
96.0
97.0
98.0
99.0
100.0

153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971

**ORIGINAL PAGE IS
OF POOR QUALITY**

A-54

ACTIVITY OF QUANTALINS, MEAN, OF STIMULATED RESPONSES TO
SERIALLY-APPLIED STIMULI
APRIL 1965



THE UNITED STATES OF AMERICA

0006-527

9901-501

15-2131


65.31967

52929

95-51276

3

1372 SORGHUM
X1-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



125.0000

105-1866

95.21311

63-11967

WS. 42623


25-57279

3

[illegible]

123 ALFA/PA/PASTJRE
K1
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION


125.0000
105.1066
85.21311
65.31367
45.42623
25.53279



0.037.....77.....117.....157.....197.....237.....

127 ALFA/PA/PASTJRE
K1
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION


125.0000
105.1066
85.21311
65.31367
45.42623
25.53279



0.037.....77.....117.....157.....197.....237.....

115 1LFA1=144V/P51JR=

141 824/EA/MA/PASTURE
X1-10-BUSINESS
-1 STANDARD DEVIATION, MEAN: 0.1 STANDARD DEVIATION



A-56

125.0300

105.1066

85.21311

65.31967


45.42323

25.5274

2.3

100


3




125.0000	105.1066	95.2111	65.31967	45.62623	25.52779
----------	----------	---------	----------	----------	----------

ORIGINAL PAGE IS
OF POOR QUALITY

9-3 ALFA/AMV/PASTJE
K1-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



9-2 ALFA/AMV/PASTJE
K1-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



125.8000

135.1366

95.21311

65.11467

-5.62623

25.53275

0.0

.....46.....47.....48.....49.....50.....51.....52.....53.....54.....55.....56.....57.....58.....59.....60.....61.....62.....63.....64.....65.....66.....67.....68.....69.....70.....71.....72.....73.....74.....75.....76.....77.....78.....79.....80.....81.....82.....83.....84.....85.....86.....87.....88.....89.....90.....91.....92.....93.....94.....95.....96.....97.....98.....99.....100.....

.....1.....2.....3.....4.....5.....6.....7.....8.....9.....10.....11.....12.....13.....14.....15.....16.....17.....18.....19.....20.....21.....22.....23.....24.....25.....26.....27.....28.....29.....30.....31.....32.....33.....34.....35.....36.....37.....38.....39.....40.....41.....42.....43.....44.....45.....46.....47.....48.....49.....50.....51.....52.....53.....54.....55.....56.....57.....58.....59.....60.....61.....62.....63.....64.....65.....66.....67.....68.....69.....70.....71.....72.....73.....74.....75.....76.....77.....78.....79.....80.....81.....82.....83.....84.....85.....86.....87.....88.....89.....90.....91.....92.....93.....94.....95.....96.....97.....98.....99.....100.....

ORIGINAL PAGE IS
OF POOR QUALITY

333 ALFALFA/HAY/PASTURE/GRASS
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

ALFALFA/HAY/PASTURE/GRASS 41-BRIGHTNESS -1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

125.0000

105.1066

85.21311


65.31967

-5.42623

25.53279

0.0

0.0 37 117 157 197 237



A-58

332 ALFALFA/HAY/PASTURE
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

ALFALFA/HAY/PASTURE 41-BRIGHTNESS -1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

125.0000

105.1066

85.21311

65.31967



-5.42623

25.53279

0.0

860 ALFALFA/44V/GRASS
41-ORIGIN:SS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

A-59



46
47
48
49
50
51
52
53
54
55

[illegible]

THE

100 ALFA/PASTURE
RI-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

865 PASTURE
K1-BRIGHTNESS
-1 STANDARD DEVIATION. MEAN. +1 STANDARD DEVIATION

125.6002

9961-501

09-21311

65.11367

MS. 62623

25.53279

9.3

1

125.0000

105-1006

05.2111

65-11467

65.4262J

25,53275

[illegible]

ORIGINAL PAGE IS
OF POOR QUALITY

A-61

002 20-SS/RY/PASTURE
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

.....

125.1200

105.1066

85.21311

65.31957

45.42623

25.53279

0.3

.....

001 ALP/PASTURE/GRASS
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

.....

125.6000

105.1066

85.21311



65.31967

45.42623

25.53279


0.0

.....



ORIGINAL PAGE IS
OF POOR QUALITY

1572 ALFALFA/PASTURE
41-BRIGHNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




0.0 37 117 157 197 237

ORIGINAL PAGE IS
OF POOR QUALITY


A-63

932 IDLE COVER
(1-REVISION)
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



932 IDLE COVER
(1-REVISION)
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

123 IDLE COVER
K1
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




123 IDLE COVER
K1
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION

ORIGINAL PAGE IS
OF POOR QUALITY


A-64

8.3 IOL COVER CRP
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



100
90
80
70
60
50
40
30
20
10
0

43. IOL COVER
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



100
90
80
70
60
50
40
30
20
10
0

125.0000

185.1065

95.21311

65.31967


65.2623

25.51273

ORIGINAL PAGE IS
OF POOR QUALITY


A-65

800 IDLE COVER
K1-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




0.0 37 77 117 157 197 237

685 IDLE COVER
K1-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION




0.0 37 77 117 157 197 237

801 IDLE COVER
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



0.037.....117.....157.....237


892 IDLE COVER
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION





0.037.....117.....157.....237

ORIGINAL PAGE IS
OF POOR QUALITY


A-66



332 FALLON FIELD
43-5K10101010
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



123 FALLON
41 -1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



0.0


0.1

0.2 0.3 0.4 0.5 0.6 0.7 0.8 0.9 1.0 1.1 1.2 1.3 1.4 1.5 1.6 1.7 1.8 1.9 2.0 2.1 2.2 2.3 2.4 2.5 2.6 2.7 2.8 2.9 3.0

0.0


ORIGINAL PAGE IS
OF POOR QUALITY

1572 FALLOM
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, .01 STANDARD DEVIATION



0.0 1.0 2.0 3.0 4.0 5.0 6.0 7.0 8.0 9.0 10.0 11.0 12.0 13.0 14.0 15.0 16.0 17.0 18.0 19.0 20.0 21.0 22.0 23.0 24.0 25.0 26.0 27.0 28.0 29.0 30.0 31.0 32.0 33.0 34.0 35.0 36.0 37.0 38.0 39.0 40.0 41.0 42.0 43.0 44.0 45.0 46.0 47.0 48.0 49.0 50.0 51.0 52.0 53.0 54.0 55.0 56.0 57.0 58.0 59.0 60.0 61.0 62.0 63.0 64.0 65.0 66.0 67.0 68.0 69.0 70.0 71.0 72.0 73.0 74.0 75.0 76.0 77.0 78.0 79.0 80.0 81.0 82.0 83.0 84.0 85.0 86.0 87.0 88.0 89.0 90.0 91.0 92.0 93.0 94.0 95.0 96.0 97.0 98.0 99.0 100.0


002 FALLOM
41-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, .01 STANDARD DEVIATION



0.0 1.0 2.0 3.0 4.0 5.0 6.0 7.0 8.0 9.0 10.0 11.0 12.0 13.0 14.0 15.0 16.0 17.0 18.0 19.0 20.0 21.0 22.0 23.0 24.0 25.0 26.0 27.0 28.0 29.0 30.0 31.0 32.0 33.0 34.0 35.0 36.0 37.0 38.0 39.0 40.0 41.0 42.0 43.0 44.0 45.0 46.0 47.0 48.0 49.0 50.0 51.0 52.0 53.0 54.0 55.0 56.0 57.0 58.0 59.0 60.0 61.0 62.0 63.0 64.0 65.0 66.0 67.0 68.0 69.0 70.0 71.0 72.0 73.0 74.0 75.0 76.0 77.0 78.0 79.0 80.0 81.0 82.0 83.0 84.0 85.0 86.0 87.0 88.0 89.0 90.0 91.0 92.0 93.0 94.0 95.0 96.0 97.0 98.0 99.0 100.0


ORIGINAL PAGE IS
OF POOR QUALITY

127 SULFLOWERS
CL-441-WINNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



ORIGINAL PAGE IS
OF POOR QUALITY

1531 CRM
K1-BRIGHTNESS
-1 STANDARD DEVIATION, MEAN, +1 STANDARD DEVIATION



125.0000

105.1366

95.21311

85.31367

63.62623

25.53279

0.0 1.0 2.0 3.0 4.0 5.0 6.0